



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

If $\cos y = .95818 + .002798 = .96098$, $y = 16^\circ 3\frac{1}{2}' = .28027^{(r)}$, a third approximation. If $\cos y = .95818 + .00281 = .96099$, $y = 16^\circ 3\frac{3}{8}' = .28023^{(r)}$, a fourth approximation. Whence $x = 96.098$, nearly.

(2). From the well known length of the side of the inscribed decagon we have $\sin 18^\circ = \cos 72^\circ = \frac{1}{4}(\sqrt{5} - 1)$.

$$\text{Hence } \cos 36^\circ = \sqrt{\frac{1 + \frac{1}{4}(\sqrt{5} - 1)}{2}} = \frac{1}{4}\sqrt{6 + 2\sqrt{5}},$$

$$\text{and } \sin 36^\circ = \sqrt{\frac{1 - \frac{1}{4}(\sqrt{5} - 1)}{2}} = \frac{1}{4}\sqrt{10 - 2\sqrt{5}}.$$

$$\text{Hence } \cot 36^\circ = \tan 54^\circ = \sqrt{\frac{6 + 2\sqrt{5}}{10 - 2\sqrt{5}}} = \sqrt{1 + \frac{2}{5}\sqrt{5}},$$

$$\tan 15^\circ = \frac{1 - \cos 30^\circ}{\sin 30^\circ} = 2 - \sqrt{3}.$$

$$\begin{aligned} \therefore \tan 39^\circ = \tan (54^\circ - 15^\circ) &= \frac{\frac{1}{\sqrt{1 + \frac{2}{5}\sqrt{5}}} - 2 + \sqrt{3}}{1 + (2 - \sqrt{3})\frac{1}{\sqrt{1 + \frac{2}{5}\sqrt{5}}}} \\ &= \frac{\sqrt{5 + 2\sqrt{5}} - 2\sqrt{5} + \sqrt{15}}{\sqrt{5} + 2\sqrt{5 + \sqrt{5}} - \sqrt{15 + 6\sqrt{5}}}. \end{aligned}$$

This can be readily computed to any desired degree of accuracy.

(3). As x varies from 0 to $\frac{1}{2}\pi$, $x^2 \tan x$ passes through all successive values from 0 to ∞ . Hence, it has one value at which $x^2 \tan x = 1$. The same is true as x varies from π to $3\pi/2$. But in the second and fourth quadrants all values of $x^2 \tan x$ are negative. Hence the equation has no real roots in those quadrants. Hence the number of real roots between 0 and 2π is 2.

Also solved by G. B. M. Zerr and A. H. Holmes. Dr. Zerr finds $\tan 39^\circ = .809785$ by means of $\tan 30^\circ$ and $\tan 9^\circ$. In (3), he finds $x = 51^\circ 17\frac{1}{2}'$ and $185^\circ 27' 10''$.

PROBLEMS FOR SOLUTION.

ALGEBRA.

279. Proposed by THEODORE L. DE LAND, Treasury Department, Washington, D. C.

The United States Panama Canal Bonds were issued, to date August 1, 1906, and will mature on August 1, 1936; and they bear interest at the rate of 2% per annum, payable quarterly, on the first day of November, 1906, and the first day of February, May, and August, 1907, and so on for each succeeding quarter, until the bonds mature, when the principal will be paid at par with the last quarter's interest. The coupon bonds of this loan were quoted on the New York Stock Exchange, at 10.30 a. m., on December 17, 1906, at 103 $\frac{3}{4}$ bid and 104 $\frac{3}{4}$ asked.

Required: The rate of interest per annum, payable quarterly, an investor would *realize* if he purchased the Panama bonds on December 17, 1906, and could reinvest his interest income, quarterly, at the *realized* rate.